



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 4, 2018

James Messina
Agent for Agrosource, Inc.
C/O Exponent LLC
1150 Connecticut Ave., NW
Suite 1100
Washington, DC 20036

Subject: PRIA Label Amendment – Oxytetracycline- New Time-Limited Uses on Citrus
Crop Group 10-10 expiring 12/4/2025
Product Name: FIRELINE 17 WP FUNGICIDE/BACTERICIDE
AGRICULTURAL OXYTETRACYCLINE
EPA Registration Number: 80990-1
Application Date: 10/23/2015
Decision Number: 510565

Dear Mr. Messina:

The application referred to above, submitted under the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable under FIFRA sec 3(c)(5) for a limited period of time. The new uses being granted in this amendment will automatically expire 12/4/2025. The following terms apply:

- 1) Resistance Management Plan implementation (e.g. education/training and stewardship plan). A yearly summary report describing the Resistance Management Plan implementation details must be submitted to the Registration Division by December 31st of each year for confirmatory purposes.
- 2) Annual sales reports listed by state, submitted by Agrosource, Inc. to the Registration Division by December 31st of each year
- 3) Monitoring requirement
 - a. You must submit protocol submissions describing how you plan to monitor both soil and citrus foliage for incidences of antibiotic resistance on a yearly basis for 3 years of this new use amendment on Citrus Crop Group 10-10. The protocol submission to the Registration Division must be made 3 months prior to each use season to allow for Agency review prior to start of monitoring. The first protocol submission is due by March 1, 2019 for the 2019 use season.
 - b. Annual monitoring report submissions by December 31st of each year starting in 2019.
- 4) One year prior to expiration, if you choose to seek an extension or remove the time limitation for the Citrus Crop Group 10-10 use, you are required to submit an application for amendment along with a revised assessment on the development of oxytetracycline resistance in human pathogens addressing release, exposure, consequence assessments

and overall risk estimation with regard to public health effects resulting from cumulative usage of oxytetracycline on all citrus crops in all states where it is registered and used. The Agency will consult with the federal partners to reevaluate the current risk picture for oxytetracycline prior to extending or removing the time limitation on citrus, or allowing the use to expire.

- 5) If the Agency determines it will not be able to grant an extension or remove the time-limitation, the Agency will notify you by 2 months prior to the use expiring. Upon notification, you must submit revised labeling with the Citrus Crop Group 10-10 use removed by 1 month prior to expiration.
- 6) You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these terms. If you have any questions, please contact Hope Johnson by phone at 703-305-5410, or via email at johnson.hope@epa.gov.

Sincerely,



Cynthia Giles-Parker, Chief
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline

OXYTETRACYCLINE	GROUP	41	FUNGICIDE
------------------------	--------------	-----------	------------------

For control or suppression of:

- Fire Blight for apples and pears
- Bacterial Spot for peaches and nectarines
- Huanglongbing (“HLB”, “citrus greening”, “greening”) for citrus Crop Group 10-10
- Citrus canker for citrus Crop Group 10-10

Active Ingredient:

Oxytetracycline Hydrochloride*	18.30%
Related Compounds	0.17%
Other Ingredients	81.53%
TOTAL	100.00%

*Equivalent to 17% oxytetracycline

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See Side/Back Panel for Additional Precautionary Statements, First Aid and Directions for Use

EPA Reg. No. 80990-1
EPA Est. No. XXXXX-XX-X

AgroSource, Inc.
P.O. Box 3091
Tequesta, Florida 33469

NOTICE: Read the entire Directions for Use and Conditions for Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product within 90 days, unopened and undamaged, and the purchase price will be refunded.

NET CONTENTS: XXXXXXXX



FIRST AID	
If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If On Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have the person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact InfoTrac at 1-800-535-5053 for emergency medical treatment information.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes or clothing. Do not breathe spray mist. Prolonged or frequent exposure may cause allergenic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco. Remove and wash contaminated clothing before reuse. This material is not to be used for medical, veterinary or human purposes.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All applicators and other handlers must wear a minimum of:

- Coveralls over short sleeved shirt and short pants
- Protective eyewear (goggles, safety glasses or face shield)
- Shoes and socks
- Chemical resistant gloves
- NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter (TC-84A); OR an elastomeric NIOSH-approved particulate respirator with any N, R or P filter (TC-84A); OR a NIOSH-approved powered air purifying respirator with an HE filter (TC-21C). Higher level respirators that are NIOSH-approved for particulates that contain oil may also be used.

Applicators must also wear chemical-resistant headgear ensuring full coverage of the neck.

Applicators do not have to wear the protective eyewear, chemical-resistant gloves, chemical-resistant headgear, or a respirator when using closed cab systems that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(e)].

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.607(d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry Interval (REI) of 12 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

Intended for use by professional applicators.

This product contains the antibiotic oxytetracycline. To reduce the development of drug-resistant bacteria and maintain the effectiveness of this and other fungicide/bactericide products, this product should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria or fungi.

Resistance Management Statements: FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline contains a Group 41 (antibiotic/fungicide/bactericide). Bacterial strains/fungal isolates with acquired resistance to Group 41 may eventually dominate the bacterial/fungal population if Group 41 products are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** or other Group 41 products.

To delay antibiotic/fungicide/bactericide resistance, take one or more of the following steps:

- Use only the specified and full strength application rates.
- The mode of action for oxytetracycline is the inhibition of protein synthesis. This product should be used to treat or prevent infections that are proven or strongly suspected to be caused by the indicated target bacteria. To reduce likelihood of bacteria developing resistance to oxytetracycline, follow the crop specific resistance management and use direction information present on this labeling. Use of this product should conform to resistance management practices/strategies established for the crop and use area (for example the use of IPM, disease forecasting models, resistance crop varieties, etc.) Consult your local extension/crop consultant or State agricultural authority if reduced efficacy is suspected.
- Adopt an integrated disease management program that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time applications.
- Avoid the consecutive use of **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** or other target site of action Group 41 products that have a similar target site of action, on the same pathogens.
- Use tank-mixtures or premixes with products from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern. Do not use any product that has a prohibition on tank mixing and follow the more restrictive use directions.
- When feasible, **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** should be alternated with a comparable bactericide with a different mode of action.
- Base use on a comprehensive IPM program.
- Monitor treated bacterial/fungal populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance contact AgroSource, Inc. at 908-931-9001.

Application Restrictions-All Uses:

- Not to be used for medical, veterinary or human purposes.
- Do not apply this product through any type of irrigation system, including chemigation.
- Do not apply this product by aerial application.
- Animal grazing in treated areas is prohibited. The public must be notified by posting restriction signs along the perimeter of the treatment area.
- Use only the specified and full-strength application rates.
- Do not apply oxytetracycline in orchards in which the soil has been fertilized with animal waste/manure or human biosolids.
- Do not apply more than two consecutive applications before alternating with another fungicide/bactericide of a different mode of action.
- Spray Drift Precaution - ALL uses - to help reduce off-target drift, direct spray into the canopy, and turn off outward pointing nozzles at row ends and when spraying outer rows.

Crop Use Directions

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
Apples	Fire Blight (<i>Erwinia amylovora</i>)	24 oz. per acre (4.4 oz. active ingredient, ai, per acre, equivalent to 4.08 oz. oxytetracycline free base)	Begin applications at the start of bloom. Repeat spray applications at 3 to 6-day intervals until the end of bloom. One additional post bloom application permitted. Apply product in sufficient water volume to achieve adequate coverage.	Maximum number of applications per year: 6 Do not apply within 60 days of harvest. Maximum <u>annual</u> amount that may be applied <u>per acre</u> : 9.0 lbs. (144 oz.) of product per acre (1.65 lbs. active ingredient per acre, equivalent to 1.53 lbs. oxytetracycline free base). Use only the full strength application rate Follow Application Restrictions-All Uses.

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
Pears	Fire Blight (<i>Erwinia amylovora</i>)	16 oz. per acre (2.93 oz. active ingredient, ai, per acre, equivalent to 2.72 oz. oxytetracycline free base)	Begin applications at 10% bloom. Repeat applications at 4 to 6-day intervals. Apply product in sufficient water volume to achieve adequate coverage.	Maximum number of applications per year: 10 Do not apply within 60 days of harvest. Maximum <u>annual</u> amount that may be applied <u>per acre</u> : 10 lbs. (160 oz.) of product per acre (1.83 lbs. active ingredient per acre,

Master Label: FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline

				<p>equivalent to 1.70 lbs. oxytetracycline free base).</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>
--	--	--	--	--

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
Peaches & Nectarines	Bacterial Spot (<i>Xanthomonas pruni</i>)	18 oz. per acre (3.29 oz. active ingredient, ai, per acre, equivalent to 3.06 oz. oxytetracycline free base)	<p>Begin applications at "shuck split". When disease pressure is high, spray at 5-7-day intervals. Consider alternating or tank mixing with copper based products. Note: certain <u>liquid</u> copper formulations may be <u>incompatible</u> with FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline.</p> <p>If disease pressure is low, spray interval may be lengthened accordingly.</p> <p>Apply product in sufficient water volume to achieve adequate coverage.</p>	<p>Maximum number of applications per year: 9</p> <p>Do not apply within 21 days of harvest.</p> <p>Maximum <u>annual</u> amount that may be applied <u>per acre</u>: 10.125 lbs. (162 oz.) of product per acre (1.85 lbs. active ingredient per acre, equivalent to 1.72 lbs. oxytetracycline free base).</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
Citrus Fruit (Crop Group 10-10) Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit;	Huanglongbing also known as "HLB", "Citrus Greening" or "Greening" disease (e.g. <i>Candidatus Liberibacter asiaticus</i> also known as	24 oz. <u>per</u> acre (4.4 oz. active ingredient, ai, per acre, equivalent to 4.08 oz. oxytetracycline free base)	<p>Apply as a foliar spray using sufficient carrier volume to ensure complete coverage. Use with spray adjuvant, spreader sticking agent or horticultural oil.</p> <p>Make first application at initiation of spring flush to suppress HLB titer and disease symptoms.</p>	<p>Maximum number of applications per year: 3</p> <p>Minimum retreatment interval: 21 days.</p> <p>Do not apply within 40 days of harvest.</p> <p>Maximum <u>annual</u> amount of product that may be applied <u>per acre</u>: 4.5 lbs. (0.82 lbs. active ingredient per acre, equivalent to</p>

Master Label: FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline

<p>Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliolate orange; uniq fruit; cultivars, varieties, and/or hybrids of these</p>	<p>"Las" or "Las")</p>		<p>Make a second application mid-summer (not less than 21 days after first application).</p> <p>Make a third application in late summer to reduce the incidence of HLB-induced fruit drop and to further suppress HLB titer and disease symptoms (not less than 21 days after second application).</p>	<p>0.76 lbs. oxytetracycline free base)</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>
---	------------------------	--	--	---

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
<p>Citrus Fruit (Citrus Crop Group 10-10) Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New</p>	<p>Citrus Canker (<i>Xanthomonas citri</i> pv. <i>citri</i>)</p>	<p>24 oz. <u>per</u> acre (4.4 oz. active ingredient, ai, per acre, equivalent to 4.08 oz. oxytetracycline free base)</p>	<p>Apply as a foliar spray using sufficient carrier volume to ensure complete coverage. Use with spray adjuvant, spreader sticking agent or horticultural oil.</p> <p>Make first application when spring flush is approximately 75% expanded to protect new growth from canker infection.</p> <p>Make a second application in summer when weather conditions warrant during periods of high citrus canker risk (not less than 21 days after first application).</p>	<p>Maximum number of applications per year: 3</p> <p>Minimum retreatment interval: 21 days.</p> <p>Do not apply within 40 days of harvest.</p> <p>Maximum <u>annual</u> amount of product that may be applied <u>per acre</u>: 4.5 lbs. (0.82 lbs. active ingredient per acre, equivalent to 0.76 lbs. oxytetracycline free base)</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>

<p>Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliolate orange; uniq fruit; cultivars, varieties, and/or hybrids of these</p>			<p>Make a third application in late summer to reduce the incidence of canker-induced fruit drop (not less than 21 days after second application).</p>	
---	--	--	--	--

MIXING: To avoid possible pesticide contamination, use only clean metal or plastic containers in preparing all solutions.

Additional information regarding use of **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** may be obtained from your local Agricultural Extension Agent or State Experimental Station.

Use of **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** may cause phytotoxicity to the fruit and/or foliage of sensitive varieties of pears, apples, peaches and nectarines and citrus (Crop Group 10-10).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep tightly closed and sealed. Product is moisture, temperature and light sensitive. Product is hygroscopic so protect from moisture. Store in a cool (<77°F, 25°C), dry place away from heat and open flames with minimum exposure to the atmosphere. Avoid extremes in temperature.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Non-refillable Container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer bag for recycling if available or dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product, which are beyond the control of AGROSOURCE, INC. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold AGROSOURCE, INC. and Seller harmless for any claims relating to such factors. AGROSOURCE, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions.

This warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or AGROSOURCE, INC., and Buyer and User assume the risk of any such use. AGROSOURCE, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. In no event shall AGROSOURCE, INC. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF AGROSOURCE, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF AGROSOURCE, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT. AGROSOURCE, INC. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of AGROSOURCE, INC.

FireLine is a trademark of AgroSource, Inc.

Manufactured in the U.S.A. by AgroSource, Inc.

©2018 AgroSource, Inc. All rights reserved

FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline Alternate Brand

Names (ABN)

Radius™ 17 WP



SUPPLEMENTAL LABELING

**FireLine™ 17 WP
Fungicide/Bactericide Agricultural Oxytetracycline**

OXYTETRACYCLINE GROUP 41 FUNGICIDE

Supplemental label to add **DIRECTIONS FOR USE** for control or suppression of:

- **Huanglongbing ("HLB", "citrus greening", "greening") for citrus Crop Group 10-10**
- **Citrus canker for citrus Crop Group 10-10**

This supplemental label expires September 24, 2019 and must not be used or distributed after this date.

Active Ingredient:

Oxytetracycline Hydrochloride*	18.30%
Related Compounds	0.17%
Other Ingredients	81.53%
TOTAL	100.00%

*Equivalent to 17% oxytetracycline

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. THIS LABELING MUST BE IN POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **FIRELINE™ 17 WP FUNGICIDE/BACTERICIDE AGRICULTURAL OXYTETRACYCLINE** BEFORE APPLYING. USE OF **FIRELINE™ 17 WP FUNGICIDE/BACTERICIDE AGRICULTURAL OXYTETRACYCLINE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **FIRELINE™ 17 WP FUNGICIDE/BACTERICIDE AGRICULTURAL OXYTETRACYCLINE (EPA REG. NO. 80990-1)**.

EPA REG. NO. 80990-1

AgroSource, Inc.
P.O. Box 3091
Tequesta, Florida 33469

A C C E P T E D
Dec 04, 2018
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 80990-1

PRECAUTIONARY STATEMENTS

The following PPE information supersedes the PPE on the label affixed to the container.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All applicators and other handlers must wear a minimum of:

- Coveralls over short sleeved shirt and short pants
- Protective eyewear (goggles, safety glasses or face shield)
- Shoes and socks
- Chemical resistant gloves
- NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter (TC-84A); OR an elastomeric NIOSH-approved particulate respirator with any N, R or P filter (TC-84A); OR a NIOSH-approved powered air purifying respirator with an HE filter (TC-21C). Higher level respirators that are NIOSH-approved for particulates that contain oil may also be used.

Applicators must also wear chemical-resistant headgear ensuring full coverage of the neck.

Applicators do not have to wear the protective eyewear, chemical-resistant gloves, chemical-resistant headgear, or a respirator when using closed cab systems that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(e)].

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.607(d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

DIRECTIONS FOR USE

Intended for use by professional applicators.

This product contains the antibiotic oxytetracycline. To reduce the development of drug-resistant bacteria and maintain the effectiveness of this and other fungicide/bactericide products, this product should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria or fungi.

Resistance Management Statements: FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline contains a Group 41 (antibiotic/fungicide/bactericide). Bacterial strains/fungal isolates with acquired resistance to Group 41 may eventually dominate the bacterial/fungal population if Group 41 products are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** or other Group 41 products.

To delay antibiotic/fungicide/bactericide resistance, take one or more of the following steps:

- Use only the specified and full strength application rates.
- The mode of action for oxytetracycline is the inhibition of protein synthesis. This product should be used to treat or prevent infections that are proven or strongly suspected to be caused by the indicated target bacteria. To reduce likelihood of bacteria developing resistance to oxytetracycline, follow the crop specific resistance management and use direction information present on this labeling. Use of this product should conform to resistance management practices/strategies established for the crop and use area (for example the use of IPM, disease forecasting models, resistance crop varieties, etc.) Consult your local extension/crop consultant or State agricultural authority if reduced efficacy is suspected.
- Adopt an integrated disease management program that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time applications.
- Avoid the consecutive use of **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** or other target site of action Group 41 products that have a similar target site of action, on the same pathogens.
- When feasible, **FireLine™ 17 WP Fungicide/Bactericide Agricultural Oxytetracycline** should be alternated with a comparable bactericide with a different mode of action.
- Base use on a comprehensive IPM program.
- For further information or to report suspected resistance contact AgroSource, Inc. at 908-931-9001.

Application Restrictions-All Uses:

- Not to be used for medical, veterinary or human purposes.
- Do not apply this product through any type of irrigation system, including chemigation.
- Do not apply this product by aerial application.
- Animal grazing in treated areas is prohibited. The public must be notified by posting restriction signs along the perimeter of the treatment area.
- Use only the specified and full-strength application rates.
- Do not apply oxytetracycline in orchards in which the soil has been fertilized with animal waste/manure or human biosolids.
- Do not apply more than two consecutive applications before alternating with another fungicide/bactericide of a different mode of action.
- Spray Drift Precaution - ALL uses - to help reduce off-target drift, direct spray into the canopy, and turn off outward pointing nozzles at row ends and when spraying outer rows.

Crop Use Directions:

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
<p>Citrus Fruit (Crop Group 10-10) Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these</p>	<p>Huanglongbing also known as "HLB", "Citrus Greening" or "Greening" disease (e.g. <i>Candidatus Liberibacter asiaticus</i> also known as "Las" or "Las")</p>	<p>24 oz. <u>per</u> acre (4.4 oz. active ingredient, ai, per acre, equivalent to 4.08 oz. oxytetracycline free base)</p>	<p>Apply as a foliar spray using sufficient carrier volume to ensure complete coverage. Use with spray adjuvant, spreader sticking agent or horticultural oil.</p> <p>Make first application at initiation of spring flush to suppress HLB titer and disease symptoms.</p> <p>Make a second application mid-summer (not less than 21 days after first application).</p> <p>Make a third application in late summer to reduce the incidence of HLB-induced fruit drop and to further suppress HLB titer and disease symptoms (not less than 21 days after second application).</p>	<p>Maximum number of applications per year: 3</p> <p>Minimum retreatment interval: 21 days.</p> <p>Do not apply within 40 days of harvest.</p> <p>Maximum <u>annual</u> amount of product that may be applied <u>per acre</u>: 4.5 lbs. (0.82 lbs. active ingredient per acre, equivalent to 0.76 lbs. oxytetracycline free base)</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>

Crop	Disease (Pathogen)	Product Application Rate	Use Directions	Specific Use Restrictions
<p>Citrus Fruit (Citrus Crop Group 10-10) Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these</p>	<p>Citrus Canker (<i>Xanthomonas citri</i> pv. <i>citri</i>)</p>	<p>24 oz. <u>per</u> acre (4.4 oz. active ingredient, ai, per acre, equivalent to 4.08 oz. oxytetracycline free base)</p>	<p>Apply as a foliar spray using sufficient carrier volume to ensure complete coverage. Use with spray adjuvant, spreader sticking agent or horticultural oil.</p> <p>Make first application when spring flush is approximately 75% expanded to protect new growth from canker infection.</p> <p>Make a second application in summer when weather conditions warrant during periods of high citrus canker risk (not less than 21 days after first application).</p> <p>Make a third application in late summer to reduce the incidence of canker-induced fruit drop (not less than 21 days after second application).</p>	<p>Maximum number of applications per year: 3</p> <p>Minimum retreatment interval: 21 days.</p> <p>Do not apply within 40 days of harvest.</p> <p>Maximum <u>annual</u> amount of product that may be applied <u>per acre</u>: 4.5 lbs. (0.82 lbs. active ingredient per acre, equivalent to 0.76 lbs. oxytetracycline free base)</p> <p>Use only the full strength application rate</p> <p>Follow Application Restrictions-All Uses.</p>

FireLine is a trademark of AgroSource, Inc.
 Manufactured in the U.S.A. by AgroSource, Inc.
 ©2018 AgroSource, Inc. All rights reserved